Lake, Fort Simpson on the Mackenzie River, Nulato on the upper Yukon, and British Columbia.

E. t. alnorum breeds in the Maritime Provinces, New England and New York, the northern part of the Lower Peninsula of Michigan (I have several typical specimens from Oden which is situated about twenty miles south of the Straits of Mackinac), and doubtless at many other points in the region lying immediately about the Great Lakes, but just how far to the westward it extends and whether or not its breeding range in the Mississippi Valley reaches sufficiently far southward to impinge on or closely approach that of true traillii, I am wholly ignorant.

# SEVENTH SUPPLEMENT TO THE AMERICAN ORNITHOLOGISTS' UNION CHECK-LIST OF NORTH AMERICAN BIRDS.

Since the publication of the Sixth Supplement to the A. O. U. Check-List the Committee on Classification and Nomenclature of North American Birds has held two sessions, having in hand the preparation of a new edition of the Check-List, as well as current questions relating to species and subspecies, and to changes in nomenclature proposed since the preparation of its last report (cf. Auk, XI, 1894, pp. 46–51). The sessions of the Committee were held in Washington. Jan. 15–18, 1894, and Feb. 12, 13, 1895, all the members of the Committee being present. In view of the proposed early publication of the second edition of the Check-List, all questions known to the Committee were considered, including nearly all of those originating in the present (April) number of 'The Auk.' Some of the latter still await final action.

The numbers at the left of the scientific names facilitate collation with the Check-List. The interpolated species and subspecies are numbered in accordance with the provision made therefor in the Code of Nomenclature (p. 14, last paragraph).

Committee Elliott Coues, Chairman.
J. A. Allen.
William Brewster.
C. Hart Merriam.
Robert Ridgway.

#### I. ADDITIONS.

### [388.1.] Cuculus canorus telephonus (Heine). Kamtschatkan Cuckoo.

Cuculus telephonus Heine, Journ. f. Orn. 1863, 352.

Cuculus canorus telephonus Stejn. Bull. 29, U. S. Nat. Mus. 1885, 224.

HAB. An Asiatic species, occurring accidentally at the Pribilof Islands. (Cf. Palmer, Auk, XI, Oct. 1894, 325.)

### [390.1.] Ceryle torquata (LINN.).

Ringed Kingfisher.

Alcedo torquata Linn. Syst. Nat. 1766, I, 180. Ceryle torquata Boie, Isis, 1828, 316.

HAB. Mexico and southward to southern South America. Casual on the Lower Rio Grande, Texas. (Cf. Stone, Auk, XI, April, 1894, 177.)

## 441.1. Basilinna leucotis (VIEILL.). White-eared Hummingbird.

Trochilus leucotis VIEILL. Ency. Méth. II, 559. Basilinna leucotis Boie, Isis, 1831, 546.

HAB. Mexico and Guatemala, north to Chiricahua Mountains, Arizona. (Cf. FISHER, Auk, XI, Oct. 1894, 325.)

480.1. Aphelocoma cyanotis RIDGW.
Blue-eared Jay.

Aphelocoma cyanotis Ridgw. Man. N. Am. Bds. 1887, 357.

HAB. Northern Mexico, ranging northward into western Texas. (Cf. Fisher, Auk, XI, Oct. 1894, 327.)

591 c. Pipilo fuscus senicula Anthony.

Anthony's Towhee.

Pipilo fuscus senicula Anthony, Auk, XII, April, 1895.

[B 396, in part, C 206 b, in part, R 240 b, in part, C 308, in part.] HAB. Southern California, and Lower California, south to Lat. 29°.

632 b. Vireo huttoni obscurus Anthony.
Dusky Vireo.

Vireo huttoni obscurus Anthony, Zoe, I, Dec. 1890, 307.

HAB. Pacific coast, from Oregon to southern British Columbia, south in winter to California.

682.1. Geothlypis poliocephala ralphi Ridgw. Rio Grande Yellow-throat.

Geothlypis poliocephala ralphi Ridgw. Proc. U. S. Nat. Mus., 1893, 692.

HAB. Lower Rio Grande Valley.

709 a. Harporhynchus cinereus mearnsi Anthony.

Mearus's Thrasher.

Harporhynchus cinereus mearnsi Anthony, Auk, XII, Jan. 1895, 53.

HAB. Northern Lower California.

713 a. Heleodytes brunneicapillus bryanti Anthony.
Bryant's Cactus Wren.

Heleodytes brunneicapillus bryanti Anthony, Auk, July. 1894, 212. [B 262, in part, C 43, in part, R 56, in part, C 63, in part.]

HAB. Northern Lower California, north into southern California.

719.1. Thryothorus leucophrys Anthony.
San Clemente Wren.

Thryothorus leucophrys Anthony, Auk, XII, Jan. 1895, 52.

HAB. San Clemente Island, California.

767. Sialia mexicana occidentalis (Towns.).
Western Bluebird.

Sialia occidentalis Towns. Journ. Acad. Nat. Sci. Phila. VII, 1837, 188.

Sialia mexicana occidentalis R1DGW. Auk, XI, April, 1894, 151, 154.

[B 159, in part, C 17, in part, R 23, in part, C 28, in part,]

HAB. Pacific Coast, from British Columbia to southern California, east to western Nevada, and casually. during migrations, to New Mexico.

767 a. Sialia mexicana bairdi RibGw.
Chestnut-backed Bluebird.

Sialia mexicana bairdi RIDGW. Auk, XI, April, 1894, 151, 157. [B 159, in part, C 17, in part, R 23, in part, C 28, in part.] HAB. Rocky Mountain district, south to northern Mexico.

767 b. Sialia mexicana anabelæ Anthony.
San Pedro Bluebird.

Sialia mexicana anabelæ Anthony, Proc. Cal. Acad. Sci., 2d Ser. II, Oct. 1889, 79. (Cf. Ridgw. Auk, XI, 1894, 151, 159.)

HAB. San Pedro Martir Mountains, Lower California.

### II. CHANGES OF NOMENCLATURE.

GENUS Engyptila SUNDEVALL (p. 180). This becomes GENUS Leptotila SWAINS.

Leptotila Swains. Class. Bds. II, 1837, 349. Type, "P. rufaxilla. Nat. Lib. V. pl. 24" [i. e., Peristera jamaicensis, Nat. Lib. IX, 207, pl. 24 (named on plate "Peristera rufaxilla")].

Leptotila Swains. has thirty-five years' priority over Engyptila Sundevall. The former name has been generally discarded as being preoccupied by Leptoptilos Vieillot, 1831, on the presumption that Swainson intended to write Leptoptila; but there being no satisfactory evidence that this was the ease, Leptotila is accepted as a tenable name.

# No. 318. Engyptila albifrons (*Bonap.*). This becomes Leptotila fulviventris brachytera (Salvad.).

Leptoptila brachyptera Salvad., Cat. Bds. Brit. Mus. XXI, 1893, 545.

Leptotila fulviventris brachyptera A. O. U. Comm. Nomen. MS.

Professor Salvadori (l. c., p. 545, footnote) has found that the type of *L. albifrons* Bonap. in the Paris Museum "is undoubtedly a specimen of *L. jamaicensis*." The earliest specific name for No. 318 of the Check-List appears therefore to be *brachyptera* Salvad. 1893 (ex Gray, 1856, nomen nudum), while *brachyptera* appears referable as a subspecies to *L. fulviventris* Lawr. (Ann. New York Acad. Sci. II, 1882, 287).

# No. 421. Chordeiles texensis Lawr. This becomes Chordeiles acutipennis texensis (Lawr.).

Chordeiles acutipennis, var. texensis Baird, Brew. & Ridgw. Hist. N. Am. Bds. II, 1874, 406. (Cf. Hartert, Cat. Bds. Brit. Mus. XVI, 1892, 616.)

Genus Picicorvus Bonap. This becomes a subgenus of Nucifraga. Cf. Ridgw. Auk, XI, April, 1894, 179.

### GENUS Nucifraga Brisson.

Nucifraga Brisson, Orn. II, 1760, 58. Type, N. caryocatactes.

Hence No. 491, Picicorvus columbianus (Wils.) becomes Nucifraga columbiana (Wils.).

Nucifraga columbiana Aud. Orn. Biog. IV, 1834, 459.

726 a. Certhia familiaris mexicana (GLOGER). This becomes

Certhia familiaris alticola MILLER.

Certhia familiaris alticola MILLER, Auk, XII, April, 1895.

#### III. ELIMINATIONS.

289 c. Colinus virginianus cubanensis (Gould). Found to be extralimital. *Cf.* Ridgway, Auk, XI, Oct. 1894, 324.

767. Sialia mexicana Swains. Found to be extralimital. Cf. Ridgway, Auk, XI, April, 1894, 151-154.

### IV. FORMS CONSIDERED AS NOT ENTITLED TO RECOGNITION.

Oceanodroma townsendi Ridgw. Proc. U. S. Nat. Mus. 1893, 687. Since the description of this species the type has been compared by Mr. Osbert Salvin with the type of *Thalassidroma melania* Bonap., in the Paris Museum, and found not to be different.

Vireo huttoni insularis Rhoads, Auk, X, July, 1893, 239. Considered as not separable from V. h. obscurus Anthony.

# V. PROPOSED CHANGES OF NOMENCLATURE REJECTED.

[253.] Totanus nebularius (Gunn.) vs. T. littorea Linn. Cf. Reichenow, J. f. O. 1889, 188. The Committee, on carefully reviewing the question, finds no reason for changing its former decision in the matter.

Bonasa umbellus fusca (ORD) COUES vs. B. u. sabini (DOUGL.). Cf. COUES, Lewis and Clark's Exped. ed. 1893, 111, 872.

It is held that the proposed change is not in accordance with Canons XLIII and XLIV of the A. O. U. Code.

### VI. ACTION DEFERRED.

Icterus gularis yucatanensis BERLEPSCH. Cf. Auk, XI, Jan. 1894, 51. Final action was again postponed.

### RECENT LITERATURE.

Newton's 'Dictionary of Birds.' Part III.1 - Part III2 of this admirable work includes, besides the minor articles to be expected in such a connection, others of considerable length and unusual importance, as naturally follows from their authoritative authorship. Among such may be especially mentioned Moa (concluded from Part II) and Odontornithes, by Mr. Lydekker; Monstrosities, Muscular System, Nervous System, and Pterylosis, by Dr. Gadow; Moor-hen, Moult, Nidification, Nightjar, Owl. Parrot, Partridge, etc., by Professor Newton. Everywhere throughout the work Professor Newton has enriched his pages with much curious historic and philological information of great interest to every intelligent reader. His views regarding nomenclature, the status and relationship of groups, and kindred general questions, are conservative and based on full knowledge of what has been said by previous authorities. He is strenuously opposed to the admission of either genera or species based on slight differences, with a tendency perhaps to underrate the fine distinctions commonly made by the writers of the 'new school.' Dr. Gadow

<sup>&</sup>lt;sup>1</sup> A | Dictionary of Birds | By | Alfred Newton | Assisted by | Hans Gadow | With Contributions from | Richard Lydekker, B. A., F. R. S., Charles S. Roy, M. A., F. R. S. | and | Robert W. Shufeldt, M. D. (late United States Army) | Part III (Moa-Sheathbill) | London | Adam and Charles Black | 1894. 8vo., pp. 577–832.

<sup>&</sup>lt;sup>2</sup> For notices of Parts I and II, see *antea*, Vol. X, pp. 357–360, and Vol. X1, pp. 56–60.